

P:\Opn\Kj\asap\p1\2378443 1 doc-5 February, 2001

Jas. A1

BIOMOLECULAR TOXICITY ASSAY

FIELD OF THE INVENTION

5 The present invention relates generally to an assay for the detection of toxicants. More particularly, the present invention contemplates an assay of toxicants such as those of the type comprising heavy metal, heavy metal divalent cations and organic molecules as well as organo-halides. Such toxicants are frequently present as contaminants in aquatic and terrestrial environments. The present invention further provides an assay device for

10 detecting toxicants. The present invention is predicated in part on the sensitivity of binding partner affinity to the toxicants.

BACKGROUND OF THE INVENTION

15 Reference to any prior art in this specification is not, and should not be taken as, an acknowledgment or any form of suggestion that this prior art forms part of the common general knowledge in Australia or any other country.

20 Bibliographic details of the publications referred to by author in this specification are collected at the end of the description.

A range of simple and complex chemical entities contaminate aquatic and terrestrial environments. Many of these molecules arise from the prolonged use of these or related or precursor molecules in a myriad of industrial processes. Such chemical entities, or

25 toxicants, include heavy metal ions, organo-halides and organic molecules. The presence of toxicants can pose serious health risks to humans and animals and can also impact on vegetation and other participants in the biosphere. Some toxicants can also have a serious impact on the ozone layer. Accordingly, the reliable analysis of toxicants is important in monitoring the environment. It is also an important component of assessing the extent of

30 contamination of industrial and urban sites which has implications for social, urban and industrial planning.

- 2 -

A range of assays have been developed for particular toxicants. Some of these assays require the use of complex and expensive apparatus and/or apparatus which is not portable or at least not readily portable. Furthermore, some assays are restricted in the range or type
5 of toxicants which can be detected. There is a need, therefore, to develop further assays for toxicants which are sensitive and inexpensive.

One type of assay which has been developed is the bioassay. Bioassays can be classified into three categories: (i) whole metazoan animal assays such as assays of the rodents, fresh
10 water copepods (crustacean) or the marine echinoderm assays (Nipper *et al.*, 1997), or plant-based assays; (ii) single cell assays such as those based on light emitted from the luminescent marine bacterium, *Photobacterium phospherum*, commonly known as the Microtox test (Ruiz *et al.*, 1997), and other single cell assays such as the lymphocyte assay (Markovic *et al.* 1977); (iii) subcellular assays using either beef heart whole mitochondria
15 or sub-mitochondria particles in which membrane-based enzyme systems are used to assay the toxicity of samples (Read *et al.*, 1997).

The range of tests currently approved by the Environmental Protection Agency (EPA) in the United States is limited to only the first of these categories. For example, the EPA has
20 nominated only eight taxa for freshwater toxicity testing and a similar number for the testing of marine toxins.

There is a need, therefore, to develop an assay which can widen the analysis base for monitoring environmental pollution. For example, in the case of heavy metal ions, the EPA
25 has set standards that described the total amount of metal ion in a sample. However, biological cellular systems separate their working biological molecules (cytoplasm) from the environment using a semi-permeable cell membrane (plasmalemma). It follows that this selectively permeable cell membrane may or may not allow a potentially polluting molecule or atom to pass through it and thus potentially toxic pollutants may or may not be
30 toxic.

Furthermore, pollutants that are toxic in some biological systems may not be toxic in other

SUMMARY OF THE INVENTION

Throughout this specification, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the
5 inclusion of a stated element or integer or group of elements or integers but not the exclusion of any other element or integer or group of elements or integers.

One aspect of the present invention provides a method for detecting an inhibitor of interaction between two or more binding partners said method comprising contacting the
10 binding partners before, during or after said partners have formed a binding partnership with a sample putatively containing said inhibitor and screening for either dissociation of binding between said binding partners or inhibition of binding between said binding partners wherein dissociation or inhibition of binding is indicative of the presence of an inhibitor.

15 Another aspect of the present invention contemplates a method for detecting an inhibitor of interaction between two or more binding partners said method comprising immobilizing a binding partner to a solid support and contacting a sample putatively containing said inhibitor before, during or after a binding partner of said first binding partner labelled with
20 a reporter molecule binds to said immobilized partner forming a binding partnership and screening for either dissociation of binding between binding partners or inhibition of binding between binding partners wherein dissociation or inhibition of binding is indicative of the presence of said inhibitor.

25 Yet another aspect of the present invention is directed to an assay device for detecting an inhibitor of interaction between members of a binding partnership comprising two or more binding partners said assay comprising a solid support comprising a binding partner immobilized thereto and a binding partner labelled with a reporter molecule to said immobilized binding partner.

30

F:\Open\A\B\...doc-5 February, 2001

Still yet another aspect of the present invention provides an assay device for detecting an inhibitor of interaction between members of a binding partnership comprising two or more binding partners said assay comprising a solid support comprising a binding partner immobilized thereto and a container adapted to contain a binding partner of said

5 immobilized binding partner which is labelled or capable of being labelled with a reporter molecule.

A further aspect of the present invention contemplates the use of binding partners of a binding partnership in the manufacture of an assay for the detection of an inhibitor of

10 interaction between said binding partners. The assay may be adapted for quantitative or quantitative deformation of the inhibitor.

This document is the property of Davies Collison Cave
 and is not to be distributed outside the organization
 without the express written permission of Davies Collison Cave
 Patent Attorneys

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154

- 6 -

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is a photographic representation showing color patterns displayed by EtBr and/or DNA applied to the nylon membrane and illuminated by UV (top panel) or an incandescent lamp (lower panel). All chemicals applied at 5 μ l volume. Additions are: (1) 10 mg/ml EtBr; (2) 0.25 mg/ml EtBr; (3) 1 mg/ml DNA; (4) DNA + 10 EtBr; (5) DNA + 0.25 EtBr at high sensitivity.

Figure 2 is a graphical representation showing changes in emission spectrum of the DNA/EtBr complex upon addition of HgCl₂. Excitation - 520 nm.

P:\Open\2\comp\pat\217111... \d\doc-3 February, 2001

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention is predicated in part on the use of macromolecular binding partners of a binding partnership in an assay to determine the presence of an inhibitor of the binding partnership.
5

Accordingly, one aspect of the present invention provides a method for detecting an inhibitor of interaction between two or more binding partners said method comprising contacting the binding partners before, during or after said partners have formed a binding partnership with a sample putatively containing said inhibitor and screening for either dissociation of binding between said binding partners or inhibition of binding between said binding partners wherein dissociation or inhibition of binding is indicative of the presence of an inhibitor.
10

Reference herein to an "inhibitor" means a molecule or chemical entity which can reduce or prevent binding between binding partners or which is capable of completely or partially dissociating binding partners after interaction. An inhibitor is preferably in the form of a toxicant. The term "toxicant" is used in its broadest sense to include any inhibitor of binding between binding partners or a molecule capable of promoting complete or partial dissociation of binding partners which have bound together in a binding partnership. Toxicants may or may not necessarily be "toxic" in the sense that they are capable of inducing death of a living organism or inducing one or more mutations in the genome of an organism or impairing physiological processes of an organism. Toxicants may also vary their toxicity depending on concentration or time of exposure. Preferred inhibitors in the form of toxicants include heavy metals, heavy metal ions and simple and complex organic molecules and organo-halides (2,4-dioxin or Picloram).
15
20
25

The sample to be tested may be a laboratory test sample, an environmental sample such as from an aquatic or terrestrial environment or may be an industrial sample from, for example, a site at or around an industrial process. A "terrestrial" sample includes gaseous samples such as air including ozone.
30

Vertical text on the left margin, possibly a page number or reference marker.

- 8 -

The assay of the present invention may be carried out in any number of ways. In one particularly useful embodiment, one member of a binding partnership is immobilized to a solid support. Another member of the binding partnership is then labelled with a reporter molecule. The binding partners are then allowed to bind. A sample putatively containing an inhibitor is then contacted with the immobilized binding partner-labelled binding partner complex. Loss of the reporter molecule is then indicative that the inhibitor has induced complete or partial dissociation of the binding partnership.

10 In another useful embodiment, the sample containing the putative inhibitor is brought into contact with the immobilized members of the binding partnership and labelled member of the binding partnership prior to binding of the partnership. Absence of the reporter molecule is indicative that the inhibitor is preventing or reducing formation of the binding partnership.

15 Generally, the binding partners represent pairs of molecules. However, the present invention extends to a binding partnership comprising three or more members.

In a particularly preferred embodiment, the binding partners are macromolecules such as proteins. Reference herein to a "protein" includes reference to a peptide or polypeptide. A protein also includes an enzyme and molecules having protein components such as glycoproteins, fusion proteins and lipoproteins. With respect to enzymes, the binding partners may include the enzymes and an enzyme substrate. Accordingly, a binding partner may also be a non-proteinaceous molecule such as a substrate. The macromolecules may also be nucleic acid molecules including RNA or DNA or hybrid forms thereof. Reference to "RNA" and "DNA" includes reference to antisense and sense molecules and oligonucleotides as well as large nucleic acid molecules. A binding partnership may also comprise nucleic acid molecules and nucleic acid binding proteins.

30 In a particularly preferred embodiment, the binding partners comprise actin and actin-binding proteins such as but not limited to cofilin and DNaseI.

- 10 -

In this particular assay, the binding partners are actin and an actin-binding protein such as cofilin or DNaseI. Globular or monomeric actin is immobilized to a solid support such as in the form of a test strip. Polymethacrylate or polystyrene are particularly preferred in this respect.

5

In practice, avidin or similar compound such as streptavidin is covalently fixed to the solid support. Actin is reacted with biotin to form biotinylated actin. The latter is then allowed to interact with the avidin/streptavidin-solid support so as to form a solid support comprising avidin/streptavidin-biotinylated actin. An actin binding partner such as cofilin or DNaseI is then labelled with a visible dye such as Texas Red (Molecular Probes, Inc.). The labelled actin binding protein is then allowed to bind to the immobilized actin and the test strip is contacted with a test sample containing, for example, a toxicant such as one or more heavy metal ions (e.g. Cd, Cu, Hg or Zn). Dissociation of the coloured binding partner is then observed when a toxicant inhibits the binding partnership. Alternatively, visualization may be following electrophoresis such as described by Kekic and dos Remedios (1999).

10
15

As stated above, the assay may be varied in any number of ways. All variations are encompassed by the present invention. For example, a solid support may comprise a wet or hydrated solid matrix where the solid support retains one binding partner that binds to a second binding partner labeled with a coloured or fluorescent ligand dye. In another alternative, the assay involves detection of a visible reaction product. In this embodiment, the toxicant alters enzymatic activity such as in whole or fragmented mitochondria. Enzyme activity is then measured, for example, spectrophotometrically to monitor absorbance changes at specific wavelengths.

20
25

Still in another alternative, an assay is based on the high affinity of toxicants such as cadmium and mercury for sulfhydryl groups which are present, for example, in many proteins. The binding to or reactive of heavy metals with sulfhydryl side chains of proteins has an effect on their reactivity.

30

In designing the assay method, it is particularly useful to select naturally-occurring

- 11 -

biological molecules such as proteins and nucleotides. However, recombinant or synthetic as well as derivative forms of these molecules may also be employed. Preferably, the binding partners interact with high affinity but be capable of interference by a range of toxins such as toxic pollutants. The term "interference" means that the interaction is altered
5 in some significant way such as a decrease in binding affinity between the binding partners. Reference to "binding" such as in a binding partnership is intended to be broadly construed and includes covalent binding including disulphide linkages, ionic bonding, hydrogen bonding and bonding or association *via* van der Waal's forces and/or electrostatic attraction.

10

The interaction between the binding partners may be detected or monitored by any means, including macroscopic and microscopic visualization. Macroscopic visualization includes visualizing light emissions and dyes as well as observing interaction between movable
solid supports such as beads.

15

The present invention further contemplates an assay device for detecting an inhibitor of
interaction between members of a binding partnership comprising two or more binding
partners said assay comprising a solid support comprising a binding partner immobilized
thereto and a binding partner labelled with a reporter molecule to said immobilized binding
20 partner.

20

In an alternative embodiment, the present invention contemplates an assay device for
detecting an inhibitor of interaction between members of a binding partnership comprising
two or more binding partners said assay comprising a solid support comprising a binding
25 partner immobilized thereto and a container adapted to contain a binding partner of said
immobilized binding partner which is labelled or capable of being labelled with a reporter
molecule.

In further embodiments, the assay device comprises containers adapted to contain agents
30 useful in conducting the assay such as reporter molecules and diluents. A container
adapted to contain a sample such as an environmental sample may also be included. The

- 12 -

assay device may be packaged separately or all together and may further comprise means to detect the presence or absence of a reporter molecule. The assay device may also contain instructions for use.

- 5 A further embodiment of the present invention contemplates the use of binding partners of a binding partnership in the manufacture of an assay for the detection of an inhibitor of interaction between said binding partners. The assay may be adapted for quantitative or qualitative determination of the inhibitor.
- 10 The term "assay" may be read synonymously with "method". Aspects of this invention relating to electrophoresis are encompassed by the description in Kekic and dos Remedios (1999) which is herein incorporated by reference.

The present invention is further described by the following non-limiting Examples.

- 13 -

EXAMPLE 1

Materials and Methods

Actin preparation, preparation of cofilin and DNaseI, native gel electrophoresis and
5 prepared or heavy metal ions and organic compounds and other methods are as described
in Kekic and dos Remedios (1999).

EXAMPLE 2

Inhibition of Interactions of Macromolecules

10

The inhibition of the interaction of, for example, actin and cofilin or actin and DNaseI can
be visualized by immobilizing one macromolecule (for example, actin) to a solid support
strip suspended in a small (1-2 ml) container. The sample is taken up into a syringe which
serves two purposes: (1) to measure the volume of the potentially polluted water sample;
15 and (2) the syringe contains a buffer compound which stabilized the pH of the aqueous
sample. A small filter is then placed on the outflow of the syringe and the sample is
injected into the vial containing the macromolecules. Dissociation of a coloured
macromolecule (for example, a Texas Red (Molecular Probes Inc.)) dye attached to cofilin
(or DNaseI or another actin-binding protein) can be visually observed to dissolve of the
20 strip containing the non-coloured actin.

The experimental basis of this assay is described below.

Four heavy metal ions (Cd, Cu, Hg and Zn) were examined over the concentration range 1-
25 1000 $\mu\text{g L}^{-1}$. The complex is most sensitive to progressive addition of 1-20 $\mu\text{g L}^{-1}$ Hg
(added as the acetate). The volume densities of the actin-cofilin bands recorded from the
native polyacrylamide gels are listed in Table 1. Addition of 1 and 5 $\mu\text{g L}^{-1}$ result in a
progressive increase in the density of the actin-cofilin complex, whereas 10 $\mu\text{g L}^{-1}$ reduces
the density of this band to levels comparable with the control with the simultaneous
30 appearance of a series of slower bands. At 20 $\mu\text{g L}^{-1}$ this trend is accentuated with the
almost complete loss of the native actin-cofilin band. The absence of unbound cofilin near

- 14 -

the top of the gel suggests that native cofilin is not dissociated. From this and other determinations. The inventors estimate that the E_{50} for Hg for the disappearance of the actin-cofilin band is approximately $20-40 \mu\text{g L}^{-1}$ using $25 \mu\text{M}$ actin-cofilin complex. This binding ratio suggests that one Hg^{2+} binds to a highly-reactive cysteinyl residue in the

5 actin-cofilin complex.

Cu^{2+} does not noticeably affect the actin-cofilin complex until it reaches a concentration of approximately $200 \mu\text{g L}^{-1}$. At higher Cu^{2+} concentrations formation of the actin-cofilin complex is progressively inhibited and the proteins seem to disperse evenly along the gel

10 lane. The 50% effective dose (E_{50}) for Cu^{2+} is approximately $400-600 \mu\text{g L}^{-1}$. Cu^{2+} can bind to a number of amino acid side chains, for example, it is known to bind to Phe-375 of the actin sequence. Judging by the E_{50} values, disruption of actin-cofilin complex is about an order of magnitude more sensitive to Hg^{2+} than to Cu^{2+} .

15 The inventors tested the effects of adding ZnCl_2 using the concentration in the range $250-2500 \mu\text{g L}^{-1}$. At these relatively high concentrations, it decreased the density of the actin-cofilin complex. The E_{50} value ($900-1000 \mu\text{g L}^{-1}$) was significantly higher than either Hg or Cu. Thus, increasing sensitivity of the actin-cofilin band to these transition metal ions was achieved in the order: $\text{Hg} \gg \text{Cu} > \text{Cd} > \text{Zn}$.

- 15 -

TABLE 1¹

Metal Cation	Actin-Cofilin µg L ⁻¹	SMP µg L ⁻¹	BHM µg L ⁻¹	Fish µg L ⁻¹ (at 96 hr)	Microtox µg L ⁻¹
Hg ²⁺	20-40	130	126	170	59
Cu ²⁺	400-600	300	93	530	9300
Cd ²⁺	800-1200	520	158	630	41400
Zn ²⁺	900-1000	1700	80	2990	33000

¹ Effective (E₅₀) concentrations for four heavy metal cations determined by: (1) Analysis of native PAGE gels of actin-cofilin; (2) Sub-mitochondrial Particle (SMP) assay; (3) Beef Heart Mitochondria (BHM) assay; (4) Whole Organism (Fish) assay; and (5) mean lethal concentrations (@ 96 hr) for *Photobacterium phosphoreum*, known as the Microtox assay. The amounts of added metal ions (g L⁻¹) did not include the mass of the counter ions (chlorides or acetate). Data for the right hand four columns were cited from Read *et al.* (1997).

EXAMPLE 3

Detection of Interaction

The actin-cofilin interaction can be monitored using a fluorescence-based detector system in which a donor fluorescent probe is excited by a light of the appropriate wavelength (using a suitable selective filter). In this example an Alexa (trade mark) fluorescent dye (used here as the donor probe) can be placed on one molecular (for example, monomeric actin) which in this case need not be fixed to a solid support. A second, different label (the acceptor probe) is reactivated with a second macromolecule (for example, cofilin), where the absorption spectrum of the acceptor is chosen to so that it overlaps the emission spectrum of the donor probe. Where there is close physical association between the actin and the cofilin an actin-cofilin complex, the emission of the donor probe is quenched by the acceptor probe and little or no fluorescence emission will occur. When the two proteins are physically dissociated (for example, a polluting toxicant), the donor emission is no longer quenched and fluorescence emission of the donor probe is detected by a photon

File: \E:\a\apples\037781...da.doc-06/02/01

detector. Thus, in this form of the device, the interaction between macromolecule ligands (for example, actin and cofilin) is monitored using fluorescence resonance energy transfer and the toxicant which dissociates the binding of these macromolecules is detected by the increase in light emission from the donor probe due to the removal of the acceptor probe.

5

EXAMPLE 4

Use of Magnetic Beads

Another form of the device relies on magnetic beads to separate the bound from the unbound proteins. Dynabeads (trade mark) provide a mechanism for sensing the binding between actin and cofilin. In this form of the device, actin is reacted with the magnetic beads and cofilin (labeled with Texas Red as above) is allowed to bind. The beads, therefore, hold actin and through actin they hold the cofilin. The proteins can be stored in this state either freeze dried or in a semi-dry state. This device works, by adding the aqueous sample to a small container containing a biological buffer. The container is fitted with a ring magnet which slides over the outside of the container and draws the magnetic beads to one end of the tube.

A water sample which failed to dissociate the coloured cofilin from the uncoloured actin is deemed safe or non-toxic. A sample which dissociated the two proteins is considered potentially toxic.

EXAMPLE 5

Nucleic Acid Sensing System

25

In this assay, a coloured indicator dye (e.g. acridine orange) is intercalated into the DNA or bound to a single-stranded nucleic acid such as RNA. Toxicity is indicated by the loss of the dye from an immobilized nucleic acid.

- 17 -

EXAMPLE 6***Electrophoresis of complexes of actin-cofilin or actin-DNaseI in native gels***

Combinations of actin, cofilin and DNaseI were mixed at equimolar concentrations (~20-
5 25 μm) in a Eppendorf tube and incubated for 15 min at 22°C. Actin, cofilin, DNaseI and
their complexes were then separated on a 10% w/v native gel. Since the separations in this
gel were achieved under nondenaturing conditions, the proteins do not migrate according
to their apparent molecular weights as they do in SDS-PAGE gels. Instead, the rate of
migration is determined by the ratio of charge to volume. Actin (M_r 43 kDa) has a net
10 negative charge. Under conditions of low ionic strength, monomeric actin is seen as a
major monomer band migrating ahead of lesser amounts of dimer and higher oligomers up
to the interface between the stacking (5%) and running (10%) gels. Cofilin is about half the
molecular mass (18.7 kDa) of actin, but because of its net positive charge at the pH
conditions in the native running gel (pH 8.6), it remains in the stacking gel. DNaseI (M_r 43
15 kDa) migrates as an indistinct band moving ahead of the actin monomers. When cofilin
and DNaseI are mixed in equimolar amounts, there is no evidence that the two form a
complex.

An equimolar complex of actin with either DNaseI or cofilin produces two bands which
20 are more sharply defined than any of the proteins alone. A mixture of actin with DNaseI
results in two sharp bands, a lower band which migrates near the actin monomer and free
DNase bands (but is not identical to either), and a slightly less dense upper band. A similar,
but nonidentical pair of bands, is seen when actin is complexed to cofilin. Note the
reduction in densities of the unbound DNaseI and cofilin bands. These actin-ABP
25 complexes are the focus of electrophoretic forms of the assays described in Kekic and dos
Remedios (1999).

- 18 -

EXAMPLE 7

Pollution test based on a protein-protein interaction where one of the proteins is immobilized on a solid matrix

- 5 In this Example, a pollution-sensitive device is described which is constructed using a solid support consisting of a wet or hydrated solid matrix where the solid support retains one protein that binds to second protein that is labeled with a colored (and or fluorescent) ligand dye and where the pollutant or toxicant inhibits the interaction of the two proteins.
- 10 The interaction of proteins like actin and cofilin has been shown to be sensitive to a number of common environmental pollutants (Kekic and dos Remedios, 1995). This Example describes a color reaction based on actin and cofilin that is easily adapted to field use and is sensitive to pollutants. The Example may also employ another actin-binding protein besides cofilin, for example, DNase I.
- 15 Cofilin is expressed as a fusion protein with a glutathione s-transferase (GST) tag. This tag allows cofilin to be bound very tightly to a matrix such as glutathione agarose which is placed in a transparent column. Actin is labeled with a highly visible fluorescent dye such as Texas Red and passed through this column. A number of these fluorescently labeled
- 20 actin molecules will remain bound to the cofilin on the column and a bright reddish color will remain that is also easily visible by illumination with ultra violet light. This coloration remains on the column after passing purified low salt buffer through it. However, the color has been shown to disappear after a solution contaminated with mercury is passed through. The removal of color is almost certainly due to the mercury ions disrupting the interaction
- 25 between the actin and cofilin. Hence, the "colored" actin is eluted from the column leaving behind the uncolored cofilin-agarose.

In practice, a polluted water sample could be detected either by the loss of coloration due to the release of a colored (labeled) second protein from the matrix or the presence of color

30 in the eluent (or both).

- 19 -

EXAMPLE 8

Pollution test based on protein activity producing a visible reaction product

An established bioassay of polluted water samples is that based on altered enzymatic
5 activity of either whole or fragmented mitochondria. Toxicants such as heavy metal ions
cause a quantitative alteration in the enzyme activities contained in these mitochondria or
mitochondrial fragments (so-called sub-mitochondrial particles). These enzymes are
usually assayed in a spectrophotometer by monitoring absorbance changes at specific
wavelengths. One of more of these mitochondrial enzymes is a suitable target for an
10 enzyme-based assay in which activity is detected by the development of a visible,
colorimetric reaction product.

EXAMPLE 9

Pollution test based on DNA-dye interaction

15 A similar device is constructed where the macromolecule is a nucleic acid like DNA bound
to a solid support (for example, but not limited to treated or untreated glass beads) retained
in a transparent tube, and where the binding partner is a dye, for example, but not limited
to fluorescent labels such as acridine orange, ethidium bromide or fluorescein-based dye.
20 Detection of a polluted water sample can be deduced by either the loss of color from the
column or by detection of the label in the column eluent.

A pollution-sensitive device is constructed, therefore, using a solid support consisting of a
wet or hydrated solid matrix where the solid support retains a nucleic acid polymer such as
25 DNA or a stabilized RNA that binds to second colored (and or fluorescent) ligand dye and
where the pollutant or toxicant inhibits the binding of the dye to the nucleic acid.

- 20 -

EXAMPLE 10***Test based on the reactivity of sulfhydryls***

Heavy metals such as cadmium and mercury have a high affinity for sulfhydryl groups
5 such as those found on many proteins. The binding to or reaction of heavy metals with
sulfhydryl side chains of proteins has an effect on their reactivity. There are a number of
established protocols for measuring the reactivity of sulfhydryls. One involves Ellmans
reagent and another method involves the use of NBD-Cl (a fluorescent dye) (Akinyele *et*
al., 1999. Another method involves the use of the fluorescent dye NBD-F. This dye is
10 reported to have a higher fluorescent yield (likely to be more visible) and a greater
reactivity (Watanabe and Imai, 1981).

Ligands may be used with any number of sulfhydryls (cysteine residues) or a cysteine
solution to detect the presence of pollutants (such as heavy metals) by measuring the effect
15 on the interaction between the fluorescent dye and the sulfhydryl. The color of the testing
solution fades visibly depending on the magnitude of the effect between the pollutant and
the sulfhydryl. This provides a field test giving a "yes/no" answer or, by using a detection
device (e.g. light transmission or absorbance) a reading is obtained - giving a high or low
value for the pollutant.

20 A pollution-sensitive device is, therefore, constructed using a solid support consisting of a
wet or hydrated solid matrix where the solid support retains a protein with multiple
cysteinyll side chains where the reactivity of these cysteinylls is reduced with pollutants or
toxicants (for example, heavy metal ions).

25

EXAMPLE 11***Test based on the protein-protein interactions using a polyacrylamide
gel-based assay using complex mixtures***

30 Heavy metals and other xenobiotics (i.e. chemicals that are foreign to life) or toxicants are
often non-toxic or minimally toxic when applied separately and individually to a

- 21 -

biotoxicity assay but can have an unexpectedly large toxicity when applied in combination (Pollak, J.K. 1998).

An electrophoretic method is used to demonstrate differential toxicity in samples of water
5 known to contain a range of pollutants.

The toxicity of combinations of chemicals (chemical mixtures) is detected by protein-protein (actin-cofilin) interactions monitored by native gel polyacrylamide gel electrophoresis or by other solution-state of solid state assay.

10

EXAMPLE 12

Immobilized DNA containing a dye/ligand is used to detect heavy metal ions

DNA is immobilized on a solid support such as but not limited to a membrane. An
15 example of this membrane is a commercial nylon membrane product called Immobilon-
Ny+ Transfer Membrane (purchased from Millipore Pty Ltd). DNA is capable of binding a
number of fluorescent dyes such as acridine orange, ethidium bromide but the dyes alone
do not bind. Sub-picogram quantities of DNA can be detected using this membrane.
Binding of a dye such as ethidium bromide (EtBr) can be detected by simple visual
20 inspection using ordinary daylight or an artificial incandescent lamp. Binding of a dye to
DNA is also detected by illuminating the DNA with ultraviolet or near-ultraviolet light and
the fluorescence visually observed (see Figure 1).

Toxicity of heavy metal ions, therefore, is detected by the inhibition of DNA-ethidium
25 bromide interaction where the DNA is immobilized on a solid membrane support.

- 22 -

EXAMPLE 13

Heavy metal ions both inhibit and alter the emission maximum of fluorescence of DNA-ligand dyes

5 The fluorescence emission of DNA-dye ligands such as ethidium bromide (EtBr) are examined spectroscopically in solution.

5 $\mu\text{g/ml}$ DNA and 2 $\mu\text{g/ml}$ EtBr at same volume of 10 μl and emission spectra are taken as presented in Figure 2. Addition of HgCl_2 resulted in concentration-dependent quenching of
10 fluorescent signal at range 1-100 μM . Changes in fluorescence intensity were noticeable at 2 μM .

The toxicity of heavy metal ions is detected, therefore, by the inhibition of DNA-ethidium bromide fluorescence where a significant decrease in fluorescence intensity can be readily
15 noted using concentrations of heavy metal ions as low as 2 μM .

EXAMPLE 14

Heavy metal ions and/or other pollutants alter conformation of proteins and thus alter the spectral properties of an attached fluorescent dye

20

When the micro-environment surrounding a fluorescent dye attached to a protein (such as actin) is altered, this results in a change to the spectral properties of this dye. These changes are either visible with the naked eye or spectrophotometrically.

25 An example of this is cadmium-mediated depolymerization of pyrene-labelled actin filaments (Wang and Templeton, 1996). This is not necessarily limited to polymeric proteins and applies to the monomeric proteins (including actin) labeled with a fluorescent dye. Such an example is the reduction in fluorescence intensity of alpha-chymotrypsinogen A labeled with 1-anilinothalene-8-sulfonic acid (ANS) when exposed to organic
30 compounds such as alcohol (Khan *et. al.*, 2000).

F:\Open\2\0\asppat\037811\037811.doc-3 February, 2001

The presence of heavy metals, organic solvents and/or other pollutants is detected, therefore, from spectral changes in proteins (or other ligands) labeled with fluorescent dyes in response to conformational and/or structural changes to the protein or ligand.

- 5 Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the invention includes all such variations and modifications. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all combinations of any two or
- 10 more of said steps or features.

The presence of heavy metals, organic solvents and/or other pollutants is detected, therefore, from spectral changes in proteins (or other ligands) labeled with fluorescent dyes in response to conformational and/or structural changes to the protein or ligand.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

- 24 -

BIBLIOGRAPHY

- Akinyele *et al.*, 1999, *J. Agric. Food Chem* 47: 2303-2307
- Kekic and dos Remedios, 1999, *Electrophoresis* 20: 2053-2058
- Khan *et al.*, 2000, *Biochim. Biophys. Acta* 1481(2): 229-236
- Markovic *et al.* 1997, *Immunol. Cell Biol.* 71 Supplement S3
- Nipper *et al.* 1997, *Ecotoxicol* 3: 109-115
- Pollak *et al.* 1998, *Int. J. Environm. Health Res.* 8: 157-163
- Read *et al.* 1997, *Environmental Applications with Submitochondrial Particles*, CRC Press, Boca Raton, pp1-35
- Ruiz *et al.* 1997, *Bull. Environ. Contam. Toxicol.* 59: 619-625
- Wang and Templeton, 1996, *Toxicol. Appl. Pharmacol.* 139(1): 115-121
- Watanabe and Imai, 1981, *Anal. Chem.* 55: 1786-1791